




L  
70  
e66  
EPT

~~K. P. SCHUBERT~~

CHECKLIST  
OF  
GEKKONOID LIZARDS



ARNOLD G. KLUGE  
Museum of Zoology  
&  
Department of Biology  
University of Michigan



SMITHSONIAN  
HERPETOLOGICAL INFORMATION  
SERVICE  
NO. 85

SMITHSONIAN  
HERPETOLOGICAL  
INFORMATION  
SERVICE

The SHIS series publishes and distributes translations, bibliographies, indices, and similar items judged useful to individuals interested in the biology of amphibians and reptiles, but unlikely to be published in the normal technical journals. Single copies are distributed free to interested individuals. Libraries, herpetological associations, and research laboratories are invited to exchange their publications with the Division of Amphibians and Reptiles.

We wish to encourage individuals to share their bibliographies, translations, etc. with other herpetologists through the SHIS series. If you have such items please contact George Zug for instructions on preparation and submission. Contributors receive 50 free copies.

Please address all requests for copies and inquiries to George Zug, Division of Amphibians and Reptiles, National Museum of Natural History, Smithsonian Institution, Washington DC 20560 USA. Please include a self-addressed mailing label with requests.



## INTRODUCTION

The following list includes all genus and species group names of extant gekkonoid lizards, their authors and years of publication. Most of the taxa I recognize are based on current common usage, and no attempt has been made to assess the monophyly of any genus. Junior objective and subjective synonyms are placed in parentheses. According to Article 80c of the ICZN, I follow existing usage in those cases under consideration by the Commission (e.g., Case 2527; BZN 46:38-40, 46-48, regarding *Nactus arnouxii* A. Duméril [1851] and *Nactus pelagicus* Girard [1858]). An asterisk is applied to all recognized subspecies, taxa followed by a question mark require further study, and a nomen nudum is followed by n.n. Not all unjustified emendations are noted in this abbreviated compilation. The list was derived from my manuscript of gekkonoid synonymies, wherein complete literature references, type localities and primary types are given. I would greatly appreciate notification of additions and corrections to the following list so that the complete version can be published with minimal errors. I wish to thank Aaron Bauer, Ross Sadlier and Glenn Shea for their assistance.

## CHECKLIST

- Aclys* Kluge, 1974  
    *concinna* Kluge, 1974 (*major*\* Storr, 1987)
- Aeluroscalabotes* Boulenger, 1887 (*Aelurosaurus* Boulenger, 1885)  
    *felinus* Günther, 1864 (*borneensis* Günther, 1864;  
        *dorsalis* W. Peters, 1871; *longicaudatus*  
        Andersson, 1924; *multituberculatus*\* [?]  
        Kopstein, 1927)
- Afroedura* Loveridge, 1944  
    *africana* Boulenger, 1888 (*namaquensis*\* FitzSimons,  
        1938; *tirasensis*\* Haacke, 1965)  
    *amatolica* Hewitt, 1925  
    *bogerti* Loveridge, 1944  
    *hawequensis* Mouton and Mostert, 1985  
    *karroica* Hewitt, 1925 (*wilmoti* Hewitt, 1926;  
        *halli*\* Hewitt, 1935)  
    *nivaria* Boulenger, 1894  
    *pondolia* Hewitt, 1925 (*multiporis*\* Hewitt, 1925;  
        *langi*\* FitzSimons, 1930; *marleyi*\* FitzSimons,  
        1930; *haackei*\* Onderstall, 1984; *major*\*  
        Onderstall, 1984)  
    *tembulica* Hewitt, 1926  
    *transvaalica* Hewitt, 1925 (*platyceps* Hewitt, 1925;  
        *loveridgei*\* Broadly, 1963)
- Agamura* Blanford, 1874 (*Rhinogekko* de Witte, 1973)  
    *femoralis* M. A. Smith, 1933

*gastropholis* Werner, 1917  
*misonnei* de Witte, 1973  
*persica* A. Duméril, 1856 (*cruralis* Blanford, 1874)

*Ailuroonyx* Fitzinger, 1843 (*Theconyx* Gray, 1845;  
*Aeluroonyx* Agassiz, 1846)  
*seychellensis* Duméril and Bibron, 1836  
*trachygaster* A. Duméril, 1851

*Alsophylax* Fitzinger, 1843 (*Altiphylax* Jeriomtschenko  
and Shcherbak, 1984)  
*laevis* Nikolsky, 1907 (*kashkarovi* Andrushko, 1968)  
*loricatus* Strauch, 1887 (*szczerbaki*\* Golubev and  
Sattarov, 1979)  
*pipiens* Pallas, 1811  
*przewalskii* Strauch, 1887 (*microtis* Blanford,  
1875)  
*tadjikiensis* Golubev, 1979  
*tokobajevi* Jeriomtschenko and Shcherbak, 1984

*Aprasia* Gray, 1839 (*Ophioseps* Barboza du Bocage, 1873;  
*Ophiopsiseps* Boulenger, 1887)  
*aurita* Kluge, 1974  
*fusca* Storr, 1979  
*haroldi* Storr, 1978  
*inaurita* Kluge, 1974  
*parapulchella* Kluge, 1974  
*pseudopulchella* Kluge, 1974  
*pulchella* Gray, 1839 (*nasutus* Barboza du Bocage,  
1873; *brevirostris* Werner, 1909)  
*repens* Fry, 1914  
*rostrata* Parker, 1956  
*smithi* Storr, 1970  
*striolata* Lütken, 1863 (*lineolata* Lütken, 1863;  
*octolineata* W. Peters, 1864; *glauerti* Parker,  
1956)

*Aristelliger* Cope, 1862 (*Idiodactylus* Bocourt, 1870;  
*Aristelligella* Noble and Klingel, 1932)  
*barbouri* Noble and Klingel, 1932  
*cochranae* Grant, 1931 (*expectatus*\* Cochran, 1933)  
*georgeensis* Bocourt, 1873 (*irregularis* Cope, 1885)  
*hechti* Schwartz and Crombie, 1975  
*lar* Cope, 1862  
*praesignis* Hallowell, 1857 (*nelsoni*\* Barbour,  
1914)

*Asaccus* Dixon and S. Anderson, 1973  
*elisae* Werner, 1895 (*eugeniae* Nikolskij, 1907;  
*ingae* Eiselt, 1973)  
*gallagheri* Arnold, 1972

- griseonotus* Dixon and S. Anderson, 1973
- Asiocolotes* Golubev, 1984  
*depressus* Minton and J. A. Anderson, 1965  
*levitoni* Golubev and Shcherbak, 1979
- Bavayia* Roux, 1913  
*crassicollis* Roux, 1913)  
*cyclura* Günther, 1872 (*neocaledonicus* Barboza du Bocage, 1873; *bavayi* Sauvage, 1878)  
*montana* Roux, 1913  
*ornata* Roux, 1913  
*sauvagii* Boulenger, 1883  
*septuiclavis* Sadlier, 1988  
*validiclavis* Sadlier, 1988
- Bogertia* Loveridge, 1941  
*lutzae* Loveridge, 1941
- Briba* Amaral, 1935  
*brasiliiana* Amaral, 1935
- Bunopus* Blanford, 1874 (*Trachydactylus* Haas and Battersby, 1959)  
*crassicauda* Nikolsky, 1907  
*spatalurus* J. Anderson, 1901 (*jolensis* Haas and Battersby, 1959; *hajarensis*\* Arnold, 1980)  
*tuberculatus* Blanford, 1874 (*blanfordii* Strauch, 1887; *gabrielis* Werner, 1936; *biporus* Werner, 1938; *abudhabi* Leviton and S. Anderson, 1967)
- Calodactylodes* Strand, 1870 (*Calodactylus* Beddome, 1870)  
*aureus* Beddome, 1870  
*illingworthi* Deraniyagala, 1953
- Carinatogekko* Golubev and Shcherbak, 1981  
*aspratilis* S. Anderson, 1973  
*heteropholis* Minton, S. Anderson, and J. A. Anderson, 1970
- Carphodactylus* Günther, 1897  
*laevis* Günther, 1897
- Chondrodactylus* W. Peters, 1870  
*angulifer* W. Peters, 1870 (*weirei* Boulenger, 1887; *namibensis*\* Haacke, 1975)
- Christinus* Wells and Wellington, 1984 (*Ridegekko* Wells and Wellington, 1985)  
*guentheri* Boulenger, 1885  
*marmoratus* Gray, 1845 (*peronii* Fitzinger, 1843; *australis* Gray, 1845; *macrodactylus* Boulenger, 1885; *affinis* Boulenger, 1885;

*biggsae* Wells and Wellington, 1985;  
*alexanderi*\* Storr, 1987)

- Cnemaspis* Strauch, 1887 (*Ancylodactylus* L. Müller, 1907; *Paragonatodes* Noble, 1921)  
*affinis* Stoliczka, 1870 (*penangensis* Flower, 1896)  
*africana* Werner, 1895  
*argus* Dring, 1978  
*barbouri* Perret, 1986  
*beddomei* Theobald, 1870 (*marmoratus* Beddome, 1870)  
*boiei* Gray, 1842  
*boulengerii* Strauch, 1887 (*glaucus* M. A. Smith, 1921)  
*dickersoni* K. P. Schmidt, 1919  
*dilepis* Perret, 1963  
*elgonensis* Loveridge, 1935  
*lavolineata* Nicholls, 1949  
*gigas* Perret, 1986  
*goaensis* Sharma, 1976  
*indica* Gray, 1846  
*jerdonii* Theobald, 1868 (*scalpensis*\* Ferguson, 1877)  
*kandiana* Kelaart, 1852 (*malabaricus* Jerdon, 1853; *gracilis* Beddome, 1870; *wicksii* Stoliczka, 1873; *humei* Theobald, 1876; *andersonii* Annandale, 1904)  
*kendalli* Gray, 1845  
*koehleri* Mertens, 1937  
*kumpoli* Taylor, 1963  
*littoralis* Jerdon, 1853 (*planiceps* Beddome, 1871)  
*mysoriensis* Jerdon, 1853  
*nairi* Inger, Marx and Koshy, 1984  
*nigridius* M. A. Smith, 1925  
*occidentalis* Angel, 1943  
*ornata* Beddome, 1870  
*petrodroma* Perret, 1986  
*podihuna* Deraniyagala, 1944  
*quattuorseriata* Sternfeld, 1912 (*bohmanni* L. Müller and Uthmoller, 1950)  
*siamensis* M. A. Smith, 1925  
*sisparensis* Theobald, 1876 (*maculatus* Beddome, 1870; *bireticulatus* Annandale, 1915)  
*spinicollis* L. Müller, 1907  
*timoriensis* Duméril and Bibron, 1836  
*tropidogaster* Boulenger, 1885  
*uzungwae* Perret, 1986  
*wynadensis* Beddome, 1870
- Coleodactylus* Parker, 1926  
*amazonicus* Andersson, 1918 (*zernyi* Wettstein, 1928)  
*brachystoma* Amaral, 1935 (*pfrimeri*

Miranda-Ribeiro, 1939)  
*guimaraesi* Vanzolini, 1957  
*meridionalis* Boulenger, 1888  
*septentrionalis* Vanzolini, 1980

*Coleonyx* Gray, 1845 (*Brachydactylus* W. Peters, 1863;  
*Anarbylus* Murphy, 1974)  
*brevis* Stejneger, 1893  
*elegans* Gray, 1845 (*scapularis* A. Duméril, 1851;  
*coleonyx* A. Duméril, 1856; *nemoralis*\*  
 Klauber, 1945)  
*fasciatus* Boulenger, 1885  
*mitratus* W. Peters, 1863 (*dovii* Boulenger, 1885)  
*reticulatus* Davis and Dixon, 1958  
*switaki* Murphy, 1974 (*gypticolus*\* Grismer and  
 Ottley, 1988)  
*variegatus* Baird, 1859 (*abbotti*\* Klauber, 1945;  
*bogerti*\* Klauber, 1945; *peninsularis*\*  
 Klauber, 1945; *slevini*\* Klauber, 1945;  
*sonoriensis*\* Klauber, 1945; *utahensis*\*  
 Klauber, 1945)

*Colopus* W. Peters, 1869  
*wahlbergii* W. Peters, 1869 (*kalaharicus*  
 FitzSimons, 1932; *furcifer*\* Haacke, 1976)

*Cosymbotus* Fitzinger, 1843 (*Platyurus* Oken, 1836;  
*Nycteridium* Günther, 1864; *Mimetozone*  
 Boulenger, 1896)  
*craspedotus* Mocquard, 1890 (*floweri* Boulenger,  
 1896)  
*platyurus* Schneider, 1792 (*schneideriana* Shaw,  
 1802; *marginatus* Cuvier, 1829; *schneideri*  
 Günther, 1864; *himalayanum* J. Anderson, 1871;  
*nepalensis* Annandale, 1907)

*Crenadactylus* Dixon and Kluge, 1964  
*ocellatus* Gray, 1845 (*bilineatus* Gray, 1845;  
*horni*\* Lucas and Frost, 1895; *naso*\* Storr,  
 1978; *rostralis*\* Storr, 1978)

*Crossobamon* Boettger, 1888 (*Ptenodactylus* Strauch, 1887)  
*eversmanni* Wiegmann, 1834 (*atropunctatus*  
 Lichtenstein and von Martens, 1856; *lumsdenii*  
 Boulenger, 1887; *maynardi*\* M. A. Smith, 1933)  
*orientalis* Blanford, 1876 (*dunstervillei* Murray,  
 1884)

*Cyrtopodion* Fitzinger, 1843 (*Dinosauria* Gistel, 1868;  
*Mesodactylus* Shcherbak and Golubev, 1984)  
*agamuroides* Nikolsky, 1900  
*elongatus* Blanford, 1875

*kachhensis* Stoliczka, 1872 (*petrensis* Murray,  
1884)  
*montiumsalsorum* Annandale, 1913  
*scaber* Heyden, 1827 (*basoglui* Baran and Gruber,  
1982)  
*watsoni* Murray, 1892 (*ingoldbyi* Procter, 1923)

*Delma* Gray, 1831 (*Nisara* Gray, 1867; *Pseudodelma* Fischer, 1882)  
*australis* Kluge, 1974  
*borea* Kluge, 1974  
*butleri* Storr, 1987 (*haroldi* Storr, 1987)  
*elegans* Kluge, 1974  
*fraseri* Gray, 1831  
*grayii* A. Smith, 1849  
*impar* Fischer, 1882 (*lineata* Rosen, 1905)  
*inornata* Kluge, 1974  
*labialis* Shea, 1987  
*mitella* Shea, 1987  
*molleri* Lütken, 1863  
*nasuta* Kluge, 1974  
*pax* Kluge, 1974  
*plebeia* De Vis, 1888 (*wollemi* [n.n.] Wells and  
Wellington, 1985)  
*tincta* De Vis, 1888 (*reticulata* Garman, 1901)  
*torquata* Kluge, 1974

*Diplodactylus* Gray, 1832 (*Stenodactylopsis*  
Steindachner, 1870; *Lucasius* Kinghorn, 1929;  
*Lucasium* Wermuth, 1965; *Turnerdactylus* Wells  
and Wellington, 1989; *Ozziedactylus* Wells and  
Wellington, 1989; *Manwellisaurus* Wells and  
Wellington, 1989)  
*alboguttatus* Werner, 1910  
*byrnei* Lucas and Frost, 1896 (*dorotheae* Wells and  
Wellington, 1985)  
*conspicillatus* Lucas and Frost, 1897 (*hillii*  
Longman, 1915; *laevis* Sternfeld, 1924;  
*platyurus* Parker, 1926)  
*damaeus* Lucas and Frost, 1896  
*fulleri* Storr, 1978  
*galeatus* Kluge, 1963  
*granariensis* Storr, 1979 (*rex\** Storr, 1988)  
*immaculatus* Storr, 1988  
*kenneallyi* Storr, 1988  
*maini* Kluge, 1962  
*mitchelli* Kluge, 1963  
*occultus* King, Braithwaite and Wombey, 1982  
*ornatus* Gray, 1845  
*polyophthalmus* Günther, 1867  
*pulcher* Steindachner, 1870 (*bilineatus* Lucas and  
Frost, 1903; *dorsotaeniata* Pellegrin, 1909;  
*dorsalis* Werner, 1910; *lucasi* Fry, 1914)



*savagei* Kluge, 1963  
*squarrosus* Kluge, 1962  
*steindachneri* Boulenger, 1885 (*stenurus* Werner,  
 1909; *jonathoni* Wells and Wellington, 1985)  
*stenodactylus* Boulenger, 1896 (*woodwardi* Fry,  
 1914)  
*taenicauda* De Vis, 1886 (*taeniocauda* Boulenger,  
 1887)  
*tessellatus* Günther, 1875 (*pachyurus* Werner, 1909)  
*vittatus* Gray, 1832 (*furcosus* W. Peters, 1864;  
*barbouri* Angel, 1936)  
*wombeyi* Storr, 1978

*Dravidogecko* M. A. Smith, 1933  
*anamallensis* Günther, 1875

*Ebenavia* Boettger, 1878  
*inunguis* Boettger, 1878 (*boettgeri* Boulenger,  
 1885)

*Eublepharis* Gray, 1827  
*angramainyu* S. Anderson and Leviton, 1966 (*ensafi*  
 Baloutch and Thireau, 1986)  
*hardwickii* Gray, 1827 (*lunatus* Blyth, 1847)  
*macularius* Blyth, 1854 (*fasciolatus\** Günther,  
 1864; *afghanicus\** Borner, 1974; *gracilis*  
 Borner, 1974; *montanus\** Borner, 1976; *fuscus\**  
 Borner, 1981; *smithi\** Borner, 1981)  
*turcmenicus* Darevsky, 1978

*Eurydactylodes* Wermuth, 1965 (*Eurydactylus* Sauvage, 1878)  
*symmetricus* Andersson, 1908  
*vieillardii* Bavay, 1869

*Geckoella* Gray, 1867  
*collegalensis* Beddome, 1870 (*speciosus* Beddome,  
 1870)  
*deccanensis* Günther, 1864 (*albofasciatus*  
 Boulenger, 1885; *deccanensis* M. A. Smith,  
 1935)  
*jeyporensis* Beddome, 1877  
*madarensis* Sharma, 1980  
*nebulosus* Beddome, 1870  
*triedrus* Günther, 1864 (*punctata* Gray, 1867)  
*yakhuna* Deraniyagala, 1945 (*zonatus* Deraniyagala,  
 1945)

*Geckolepis* Grandidier, 1867  
*anomala* Mocquard, 1902  
*maculata* W. Peters, 1880 (*humbloti* Vaillant, 1887)  
*modesta* Methuen and Hewitt, 1913  
*petiti* Angel, 1942

*polylepis* Boettger, 1893  
*typica* Grandidier, 1867

*Geckonia* Mocquard, 1895

*chazaliae* Mocquard, 1895 (*malazodes* Ahl, 1930)

*Gehyra* Gray, 1834 (*Peropus* Wiegmann, 1835; *Phyria* Gray, 1842; *Dactyloperus* Fitzinger, 1843; *Perodactylus* Fitzinger, 1843; *Peripia* Gray, 1845; *Chalinocnemis* Duges, 1883; *Spasmocnemis* Duges, 1883; *Phyriadora* Wells and Wellington, 1985)

*angusticaudata* Taylor, 1963

*australis* Gray, 1845 (*punctulata* Gray, 1842)

*baliola* A. Duméril, 1851 (*marmorata* Macleay, 1877; *brevicaudis* Macleay, 1877)

*barea* Kopstein, 1926

*borroloola* King, 1983

*brevipalmata* W. Peters, 1874

*butleri* Boulenger, 1900

*catenata* Low, 1979

*dubia* Macleay, 1877 (*longicaudis* Macleay, 1877; *torresiana* Günther, 1877)

*fehlmanni* Taylor, 1962

*intermedia* Brown, 1902

*interstitialis* Oudemans, 1894

*kimberleyi* Borner and Schuttler, 1983

*lacerata* Taylor, 1962

*lampei* Andersson, 1913

*leopoldi* Brongersma, 1930

*marginata* Boulenger, 1887 (*fischeri* Strauch, 1887; *marginata* Boettger, 1895)

*membranacruralis* King and Horner, 1989

*minuta* King, 1982

*montium* Storr, 1982

*mutilata* Wiegmann, 1835 (*peronii* Duméril and Bibron, 1836; *platurus* Bleeker, 1858;

*insulensis* Girard, 1858; *pardus* Tytler, 1865;

*perdicolor* Cope, 1868; *packardii* Cope, 1869;

*navarri* Duges, 1883; *beebei* Annandale, 1913)

*nana* Storr, 1978

*occidentalis* King, 1984

*oceanica* Lesson, 1830 (*pacifica* Gray, 1834;

*oualensis* Duméril and Bibron, 1836; *gularis* Gray, 1842)

*pamela* King, 1982

*papuana* Meyer, 1874

*pilbara* Mitchell, 1965 (*cognata* Borner and Schuttler, 1982)

*punctata* Fry, 1914 (*fenestra* Mitchell, 1965)

*purpurascens* Storr, 1982

*robusta* King, 1983

*variegata* Duméril and Bibron, 1836 (*grayi* Steindachner, 1867; *pusillus* [?] Cope, 1868; *mestoni* de Vis, 1890; *annetteae* Wells and Wellington, 1985; *kingi* Wells and Wellington, 1985; *lazelli* Wells and Wellington, 1985)  
*vorax* Girard, 1858  
*xenopus* Storr, 1978

*Gekko Laurenti*, 1768 (*Gecko* Brongniart, 1800; *Gecus* Rafinesque, 1810; *Platydyctylus* Goldfuss, 1820; *Gecco* Ritgen, 1828; *Lomatodactylus* van der Hoeven, 1833; *Scelotretus* Fitzinger, 1843)

*athymus* Brown and Alcalá, 1962  
*auriverrucosus* Zhou and Liu, 1982  
*chinensis* Gray, 1842 (*similignum* M. A. Smith, 1923)

*gecko* Linnaeus, 1758 (*verticillatus* Laurenti, 1768; *teres* Laurenti, 1768; *guttatus* Daudin, 1802; *verus* Merrem, 1820; *annulatus* Kuhl, 1820; *reevesii* Gray, 1831; *tenuis* Hallowell, 1857; *indicus* Girard, 1858; *azhari*\* Mertens, 1955)

*gigante* Brown and Alcalá, 1978  
*hokouensis* Pope, 1928 (*amissus* Taylor, 1962)  
*japonicus* Schlegel (*in* Duméril and Bibron), 1836 (*jamori* Temminck and Schlegel, 1838; *nanus* Cantor, 1842; *yamori* Fritze, 1891)

*kikuchii* Oshima, 1912  
*liboensis* Zhou and Li, 1982  
*mindorensis* Taylor, 1919  
*palawanensis* Taylor, 1925  
*palmatus* Boulenger, 1907  
*petricolus* Taylor, 1962  
*porosus* Taylor, 1922

*romblon* Brown and Alcalá, 1978  
*scabridus* Liu and Zhou, 1982  
*smithii* Gray, 1842 (*stentor* Cantor, 1847; *albomaculatus* Giebel, 1861; *albofasciolatus* Günther, 1867)

*subpalmatus* Günther, 1864 (*melli* Vogt, 1922)

*swinhonis* Günther, 1864  
*taibaiensis* Mingtao, 1985  
*tawaensis* Okada, 1956

*tuberculatus* Daudin, 1802 (*monarchus* Schlegel, *in* Duméril and Bibron, 1836; *burmeisteri* Giebel, 1861; *deissneri* Giebel, 1861)

*verreauxii* Tytler, 1865  
*vittatus* Houttuyn, 1782 (*unistriata* Shaw, 1792; *bivittatus* Duméril and Bibron, 1836; *trachyloemus* W. Peters, 1872)

*yakuensis* Matsui and Okada, 1968

- Gonatodes* Fitzinger, 1843  
*albogularis* Duméril and Bibron, 1836 (*fuscus\** Hallowell, 1855; *varius* A. Duméril, 1856; *notatus\** Reinhardt and Lütken, 1862; *maculatus* Steindachner, 1867; *braconnieri* O'Shaughnessy, 1875; *bodinii\** Rivero Blanco, 1964)  
*annularis* Boulenger, 1887 (*boonii* Lidth de Jeude, 1904; *beebei* Noble, 1923)  
*antillensis* Lidth de Jeude, 1887  
*atricucullaris* Noble, 1921  
*caudiscutatus* Günther, 1859 (*collaris* Garman, 1892)  
*ceciliae* Donoso-Barros, 1965  
*concinatus* O'Shaughnessy, 1881 (*buckleyi* O'Shaughnessy, 1881; *ligiae* Donoso-Barros, 1967)  
*eladioi* Nascimento, Avila-Pires and Cunha, 1987  
*falconensis* Shreve, 1947  
*hasemani* Griffen, 1917 (*spinulosus* Amaral, 1932)  
*humeralis* Guichenot, 1855 (*ferriugineus* 24Cope, 1864; *incertus* W. Peters, 1871; *sulcatus* O'Shaughnessy, 1875)  
*ocellatus* Gray, 1831 (*confidentalis* Marcuzzi, 1950)  
*petersi* Donoso-Barros, 1967  
*seigliei* Donoso-Barros, 1965  
*taniae* Roze, 1963  
*tapajonicus* Rodrigues, 1980  
*vittatus* Wiegmann, 1856 (*gilli* Cope, 1864; *roquensis\** Roze, 1956)
- Goniurosaurus* Barbour, 1908 (*Amamisaurus* Borner, 1981)  
*kuroiwae* Namiye, 1912 (*orientalis\** Maki, 1931; *yamashinae* Okada, 1936; *splendens\** Nakamura and Ueno, 1959)  
*lichtenfelderi* Mocquard, 1897 (*hainanensis\** Barbour, 1908)
- Gonydactylus* Kuhl and van Hasselt, 1822 (*Goniodactylus* Schlegel, 1826; *Cyrtodactylus* Gray, 1827; *Gonyodactylus* Wagler, 1830; *Goniodactylus* Gray, 1842; *Gonyodactylus* Fitzinger, 1843; *Puellula* Blyth, 1860; *Quantasia* Wells and Wellington, 1985)  
*agusanensis* Taylor, 1915  
*angularis* M. A. Smith, 1921  
*annulatus* Taylor, 1915  
*baluensis* Mocquard, 1890  
*biordinis* Brown and McCoy, 1980  
*brevipalmatus* M. A. Smith, 1923

*cavernicolus* Inger and King, 1961 (*kinabalensis*  
Malkmus, 1989)  
*chitralensis* M. A. Smith, 1935  
*condorensis* M. A. Smith, 1921  
*consobrinoides* Annandale, 1905  
*consobrinus* W. Peters, 1871  
*darmandvillei* Weber, 1890 (*defossei* Dunn, 1927)  
*dattanensis* Khan, 1980  
*derongo* Brown and Parker, 1973  
*deveti* Brongersma, 1948  
*elok* Dring, 1978  
*fasciolatus* Blyth, 1860  
*feae* Boulenger, 1893  
*fraenatus* Günther, 1864  
*fumosus* F. Müller, 1895 (*halmahericus\** Mertens,  
1929)  
*gubernatoris* Annandale, 1913  
*himalayansis* Duda and Sahi, 1978  
*indusoani* Khan, 1988  
*ingeri* Hikida, 1990  
*intermedius* M. A. Smith, 1917  
*irregularis* M. A. Smith, 1921  
*jellesmae* Boulenger, 1897  
*khasiensis* Jerdon, 1870 (*himalayicus* Annandale,  
1906; *tamaiensis\** M. A. Smith, 1940)  
*kirmanensis* Nikolsky, 1900  
*laevigatus* Darevsky, 1964 (*uniformis\** Auffenberg,  
1980)  
*lateralis* Werner, 1896  
*lawderanus* Stoliczka, 1871 (*himalayensis*  
Annandale, 1913)  
*loriae* Boulenger, 1897  
*lousiadensis* de Vis, 1892 (*tuberculatus* Lucas and  
Frost, 1900; *olivii* Garman, 1901)  
*malcolmsmithi* Constable, 1949  
*malayanus* de Rooij, 1915  
*mansarulus* Duda and Sahi, 1978  
*marmoratus* Gray, 1831 (*agamensis* Bleeker, 1860)  
*matsuii* Hikida, 1990  
*mimikanus* Boulenger, 1914  
*mintoni* Golubev and Shcherbak, 1981  
*novaeguineae* Schlegel, 1844  
*oldhami* Theobald, 1876  
*papuensis* Brongersma, 1928 (*novaeguineae*  
Brongersma, 1928)  
*peguensis* Boulenger, 1893 (*zebraicus\** Taylor,  
1962)  
*philippinicus* Steindachner, 1867  
*pubisulcus* Inger, 1957  
*pulchella* Gray, 1827  
*quadrivirgatus* Taylor, 1962  
*rohtasfortai* Khan and Tasnim, 1990

*redimiculus* King, 1962  
*rubidus* Blyth, 1860 (*tigris* Tytler, 1864)  
*sadleiri* Wells and Wellington, 1985  
*sermowaiensis* de Rooij, 1915  
*stoliczkai* Steindachner, 1867 (*yarkandensis* J. Anderson, 1872; *walli* Ingoldby, 1922)  
*sworderi* M. A. Smith, 1925  
*tibetinus* Boulenger, 1905  
*variegatus* Blyth, 1859  
*wetariensis* Dunn, 1927  
*yoshii* Hikida, 1990

*Gymnodactylus* Spix, 1825 (*Dasyderma* Fitzinger, 1843)  
*geckoides* Spix, 1825 (*spixii* Gray, 1831;  
*gymnodactylus* Schinz, 1833; *spinulosus*  
 Fitzinger, 1843; *darwinii*\* Gray, 1845;  
*girardi* Steindachner, 1867; *amarali*\* Barbour,  
 1925; *conspicuus* Amaral, 1932; *helgae* Amaral,  
 1950)  
*guttulatus* Vanzolini, 1982

*Hemidactylus* Gray, 1825 (*Gecus* Rafinesque, 1810;  
*Hemidact* Oken, 1817; *Boltalia* Gray, 1842;  
*Hoplopodion* Fitzinger, 1843; *Microdactylus*  
 Fitzinger, 1843; *Onychopus* Fitzinger, 1843;  
*Tachybates* Fitzinger, 1843; *Pnoepus*  
 Fitzinger, 1843; *Doryura* Gray, 1845;  
*Velernesia* Gray, 1845; *Leiurus* Gray, 1845;  
*Nubilia* Gray, 1845; *Eurhous* Fitzinger, 1861;  
*Liurus* Cope, 1862; *Emydactylus* Bocourt, 1870;  
*Bunocnemis* Günther, 1894; *Lophopholis* M. A.  
 Smith and Deraniyagala, 1934; *Aliurus* Dunn  
 and Dunn, 1940)  
*agrius* Vanzolini, 1978  
*albopunctatus* Loveridge, 1942  
*aporus* Boulenger, 1906  
*arnoldi* Lanza, 1978  
*barodanus* Boulenger, 1901  
*bavazzanoi* Lanza, 1978  
*bayonii* Barboza du Bocage, 1893  
*bouvieri* Bocourt, 1870 (*cessacii* Barboza du  
 Bocage, 1873; *boavistensis*\* Boulenger, 1906;  
*chevalieri* Angel, 1935; *razoensis*\* Gruber and  
 Schleich, 1982)  
*bowringii* Gray, 1845 (*berdmorei* Blyth, 1853)  
*brookii* Gray, 1845 (*angulatus*\* Hallowell, 1854;  
*tytleri* Tytler, 1865; *guineensis* W. Peters,  
 1868; *affinis* Steindachner, 1870;  
*kushmorensis* Murray, 1884; *gleadowi* Murray,  
 1884; *stellatus* Boulenger, 1885; *murrayi*  
 Gleadow, 1887; *tenkatei* Lidth de Jeude, 1895;  
*togoensis* Werner, 1897; *haitianus*\* Meerwarth,

- 1901; *subtriedroides*\* Annandale, 1905;  
*leightoni*\* Boulenger, 1911; *luzonensis*  
 Taylor, 1915; *neotropicalis* Shreve, 1936;  
*parvimaculatus*\* Deraniyagala, 1953)  
*citernii* Boulenger, 1912  
*curlei* Parker, 1942  
*depressus* Gray, 1842 (*argentii* Gray, 1845;  
*pieressii* Kelaart, 1853)  
*echinus* O'Shaughnessy, 1875  
*fasciatus* Gray, 1842 (*ornatus* Gray, 1845; *formosus*  
 Hallowell, 1857; *ituriensis*\* K. P. Schmidt,  
 1919)  
*flaviviridis* Rüppell, 1835 (*coctaei* Duméril and  
 Bibron, 1836; *sublevis* Gray, 1842; *cocteau*  
 Fitzinger, 1843; *ruppelli* Fitzinger, 1843;  
*bengaliensis* J. Anderson, 1871; *sublaevis*  
 Boulenger, 1885; *bengalensis* Boulenger, 1885;  
*zollii* Scortecci, 1929)  
*forbesii* Boulenger, 1899  
*frenatus* Duméril and Bibron, 1836 (*bojeri*  
 Fitzinger, 1843; *javanicus* Fitzinger, 1843;  
*vittatus* Gray, 1845; *punctatus* Jerdon, 1853;  
*inornatus* Hallowell, 1861; *pumilus* Hallowell,  
 1861; *caracal* Tytler, 1865; *chaus* Tytler,  
 1865; *hexaspis* Cope, 1869; *longiceps* Cope,  
 1869; *papuensis* Macleay, 1877; *tristis*  
 Sauvage, 1878; *fragilis* Calabresi, 1915;  
*vandermeermohri* Brongersma, 1928; *okinawensis*  
 Okada, 1936)  
*funaiolii* Lanza, 1978  
*garnotii* Duméril and Bibron, 1836 (*peruvianus*  
 Wiegmann, 1835; *vulpecula* Girard, 1858;  
*ludekingii* Bleeker, 1858; *guadama* Theobald,  
 1868; *mortoni* Theobald, 1868; *mandelianus*  
 Stoliczka, 1871; *blanfordii* Boulenger, 1885)  
*giganteus* Stoliczka, 1871  
*gracilis* Blanford, 1870 (*platyceps* Annandale,  
 1912)  
*granchii* Lanza, 1978  
*granti* Boulenger, 1899  
*greefi* Barboza du Bocage, 1886  
*homoeolepis* Blanford, 1881  
*intestinalis* Werner, 1897 (*ansorgii* Boulenger,  
 1901)  
*isolepis* Boulenger, 1895  
*jubensis* Boulenger, 1895  
*karenorum* Theobald, 1868  
*klauberi* Scortecci, 1948  
*laevis* Boulenger, 1901  
*laticaudatus* Andersson, 1910 (*fossatii*\* Scortecci,  
 1928)  
*lemurinus* Arnold, 1980

- leschenaultii* Duméril and Bibron, 1836 (*bellii* Gray, 1845; *pustulosus* Lichtenstein and von Martens, 1856; *kelaarti* Theobald, 1868; *marmoratus* Blanford, 1870)  
*longicephalus* Barboza du Bocage, 1873 (*longiceps* O'Shaughnessy, 1873; *bocagii* Boulenger, 1885; *molleri* Bedriaga, 1892; *hecqui* Boulenger, 1901; *steindachneri* Tornier, 1902)  
*mabouia* Moreau de Jonnès, 1818 (*tuberculosus* Raddi, 1823; *incanescens* Wied-Neuwied, 1824; *armatus* Wied-Neuwied, 1824; *aculeatus* Spix, 1825; *cruciger* Spix, 1825; *mabuia* Cuvier, 1829; *mabuya* Fitzinger, 1843; *calabaricus* Boettger, 1878; *benguellensis* Barboza du Bocage, 1893)  
*macropholis* Boulenger, 1896  
*maculatus* Duméril and Bibron, 1836 (*sykesii* Günther, 1864; *hunae\** Deraniyagala, 1937)  
*marmoratus* Hallowell, 1861  
*matschiei* Toriner, 1901  
*megalops* Parker, 1932  
*mercatorius* Gray, 1842 (*gardineri* Boulenger, 1909; *persimilis* Barbour, 1928; *mandanus* Loveridge, 1936)  
*modestus* Günther, 1894  
*muriceus* W. Peters, 1870  
*newtoni* Ferreira, 1897  
*nigriventris* Lidth de Jeude, 1905  
*ophiolepis* Boulenger, 1903  
*ophiolepidoides* Lanza, 1978  
*oxyrhinus* Boulenger, 1899  
*palaichthus* Kluge, 1969  
*persicus* J. Anderson, 1872 (*bornmuelleri* Werner, 1895)  
*platycephalus* W. Peters, 1854  
*prashadi* M. A. Smith, 1935  
*puccionii* Calabresi, 1927  
*pumilio* Boulenger, 1899 (*pumilus* Boulenger, 1899)  
*reticulatus* Beddome, 1870  
*richardsonii* Gray, 1845  
*ruspolii* Boulenger, 1896 (*erlangeri* Steindachner, 1907)  
*scabriceps* Annandale, 1906  
*sinaitus* Boulenger, 1885 (*shugraensis* Haas and Battersby, 1959)  
*smithi* Boulenger, 1895  
*somalicus* Parker, 1932  
*squamulatus* Tornier, 1896 (*bocagei* Toriner, 1896; *wernerii* Tornier, 1897; *tornieri* Mocquard, 1902; *floweri\** Werner, 1908; *alluandi* Angel, 1923; *barbouri\** Loveridge, 1942)  
*stejnegeri* Ota and Hikida, 1989



*subtriedrus* Jerdon, 1853  
*tanganicus* Loveridge, 1929  
*tasmani* Hewitt, 1932  
*taylori* Parker, 1932  
*triedrus* Daudin, 1802 (*lankae*\* Deraniyagala, 1953)  
*tropidolepis* Mocquard, 1888  
*turcicus* Linnaeus, 1758 (*cyanodactylus* Rafinesque, 1810; *meridionalis* Risso, 1826; *granosus* Heyden, 1827; *robustus* Heyden, 1827; *veruculatus* Cuvier, 1829; *verrucosus* Gray, 1831; *karachiensis* Murray, 1884; *exsul* Barbour and Cole, 1906; *parkeri*\* Loveridge, 1936; *spinalis*\* Buchholz, 1954)  
*vietnamensis* Darevsky and Kupriyanova, 1984  
*yerburii* J. Anderson, 1895 (*pauciporosus*\* Lanza, 1978)

*Hemiphyllodactylus* Bleeker, 1860 (*Spathodactylus* Günther, 1872; *Spathoscalabotes* Boulenger, 1885; *Cainodactylus* Barbour, 1924)  
*harterti* Werner, 1900 (*larutensis* Boulenger, 1900)  
*insularis* Taylor, 1918  
*typus* Bleeker, 1860 (*gracilis* Bleeker, 1860; *crepuscularis* Bavay, 1869; *aurantiacus*\* Beddome, 1870; *mutilatus* Günther, 1872; *ceylonensis* Boulenger, 1885; *leucostictus* Stejneger, 1899; *margarethae* Brongersma, 1931; *chapaensis*\* Bourret, 1937; *pallidus*\* Auffenberg, 1980)  
*yunnanensis* Boulenger, 1903 (*dushanensis*\* Zhou and Liu, 1981; *jinpingsensis*\* Zhou and Liu, 1981; *longlingensis*\* Zhou and Liu, 1981)

*Hemitheconyx* Stejneger, 1893 (*Psilodactylus* Gray, 1864)  
*caudicinctus* A. Duméril, 1851  
*taylori* Parker, 1930

*Heteronotia* Wermuth, 1965 (*Heteronota* Gray, 1845)  
*binoei* Gray, 1845 (*derbianus* Gray, 1845; *australis* Steindachner, 1867; *anomalus* W. Peters, 1868; *annulatus* Macleay, 1877; *horneri* Wells and Wellington, 1985; *wadei* Wells and Wellington, 1985)  
*planiceps* Storr, 1989  
*spelea* Kluge, 1963

*Holodactylus* Boettger, 1893  
*africanus* Boettger, 1893 (*aculeatus* Calabresi, 1915)  
*cornii* Scortecci, 1931

- Homonota* Gray, 1845 (*Cubina* Gray, 1845; *Cubinia* Gray, 1845; *Wallsaurus* Underwood, 1954; *Garthia* Donoso-Barros and Vanzolini, 1965; *Garthia* Vanzolini and Donoso-Barros, 1966)
- andicola* Cei, 1978
- borellii* Peracca, 1897
- darwinii* Boulenger, 1885 (*macrocephala*\* Cei, 1978)
- fasciatus* Duméril and Bibron, 1836 (*pasteuri* Wermuth, 1965)
- gaudichaudii* Duméril and Bibron, 1836 (*dorbignii*\* Duméril and Bibron, 1836; *klugei*\* Donoso-Barros, 1970)
- horrida* Burmeister, 1861 (*mattogrossensis* Berg, 1895)
- penai* Donoso-Barros, 1965
- underwoodi* Kluge, 1964
- uruguayensis* Vaz-Ferreira and Soriano, 1961
- whitii* Boulenger, 1885
- Homopholis* Boulenger, 1885 (*Platypholis* Boulenger, 1890; *Blaesodactylus* Boettger, 1893)
- antongilensis* Bohme and Meier, 1980
- boivini* A. Duméril, 1856
- fasciata* Boulenger, 1890 (*erlangeri*\* Steindachner, 1906)
- mulleri* Visser, 1987
- sakalava* Grandidier, 1867 (*heterolepis* Boulenger, 1896)
- walbergii* A. Smith, 1849 (*macrolepis* Boulenger, 1885; *arnoldi* Loveridge, 1944)
- Hoplodactylus* Fitzinger, 1843 (*Pentadactylus* Gray, 1845; *Dactylocnemis* Steindachner, 1867; *Woodworthia* Garman, 1901)
- chrysosireticus* Robb, 1980
- delcourti* Bauer and Russell, 1986
- duvaucelii* Duméril and Bibron, 1836
- granulatus* Gray, 1845 (*brevidactylus* Gray, 1845; *greyii* Knox, 1870; *sylvestris* Buller, 1881; *versicolor* Colenso, 1885; *nebulosus* McCann, 1955)
- kahutarae* Whitaker, 1985
- maculatus* Boulenger, 1885
- pacificus* Gray, 1842 (*pomarii* Girard, 1858; *wullerstorfi* Fitzinger, 1861; *brunneus* Cope, 1868; *digitatus* Garman, 1901)
- rakiurae* Thomas, 1981
- stephensi* Robb, 1980
- Kaokogecko* Steyn and Haacke, 1966
- vanzyl* Steyn and Haacke, 1966

- Lepidoblepharis* Peracca, 1897 (*Lathrogecko* Ruthven, 1916)  
*buchwaldi* Werner, 1910  
*colombianus* Mechler, 1968  
*duolepis* Ayala and Castro, 1983  
*festae* Peracca, 1897  
*grandis* Miyata, 1985  
*heyerorum* Vanzolini, 1978  
*intermedius* Boulenger, 1914  
*miyata* Lamar, 1985  
*microlepis* Noble, 1923  
*oxycephalus* Werner, 1894  
*peraccae* Boulenger, 1908  
*ruthveni* Parker, 1926  
*sanctaemartae* Ruthven, 1916 (*fugax* Ruthven, 1928)  
*williamsi* Ayala and Serna, 1986  
*xanthostigma* Noble, 1916
- Lepidodactylus* Fitzinger, 1843 (*Amydosaurus* Gray, 1845)  
*aureolineatus* Taylor, 1915  
*balioburius* Ota and Crombie, 1989  
*browni* Pernetta and Black, 1983  
*christiani* Taylor, 1917  
*euaensis* Gibbons and Brown, 1988  
*gardineri* Boulenger, 1897  
*guppyi* Boulenger, 1884  
*herrei* Taylor, 1923 (*medianus*\* Brown and Alcalá, 1978)  
*listeri* Boulenger, 1889  
*lugubris* Duméril and Bibron, 1836 (*neglectus* Girard, 1858; *meijeri* Bleeker, 1859; *pomareae* Fitzinger, 1861; *cantoris* Günther, 1864; *harrieti* Tytler, 1865; *moestus* W. Peters, 1867; *roseus* Cope, 1869; *caudeloti* Bavay, 1869; *mysorensis* Meyer, 1874; *ornata* Macleay, 1877; *divergens* Taylor, 1918; *lombocensis* Mertens, 1929; *ogasawarasimae* Okada, 1930; *intermedius*\* Darevsky, 1964)  
*magnus* Brown and Parker, 1977  
*manni* K. P. Schmidt, 1923  
*mutahi* Brown and Parker, 1977  
*novaeguineae* Brown and Parker, 1977  
*oorti* Kopstein, 1926  
*orientalis* Brown and Parker, 1977  
*planicaudus* Stejneger, 1905 (*naujanensis* Taylor, 1919)  
*pulcher* Boulenger, 1885  
*pumilus* Boulenger, 1885  
*pusillus* [?] Cope, 1868  
*ranauensis* Ota and Hikida, 1988  
*shebae* Brown and Tanner, 1949  
*woodfordi* Boulenger, 1887

yami Ota, 1987

*Lialis* Gray, 1835 (*Ophiophthalmus* Fitzinger, 1843;  
*Alopecosaurus* Lindholm, 1905)  
*burtonis* Gray, 1835 (*bicatenata* Gray, 1842;  
*punctulata* Gray, 1842; *leptorhyncha* W.  
 Peters, 1874; *concolor* W. Peters, 1874)  
*jicari* Boulenger, 1903 (*cuneirostris* Lindholm,  
 1905; *inornata* Lindholm, 1905)

*Luperosaurus* Gray, 1845  
*brooksii* Boulenger, 1920  
*browni* Russell, 1979  
*cumingii* Gray, 1845  
*joloensis* Taylor, 1918  
*macgregori* Stejneger, 1907  
*palawanensis* Brown and Alcalá, 1974

*Lygodactylus* Gray, 1864 (*Scalabotes* W. Peters, 1880;  
*Vanzoia* H. M. Smith, Martin and Swain, 1977)  
*angolensis* Barboza du Bocage, 1896 [1897] (*laurae*  
 K. P. Schmidt, 1933)  
*angularis* Günther, 1893 (*heeneeni*\* DeWitte, 1933;  
*grzimeki*\* Bannikov and Darevsky, 1969)  
*arnoulti* Pasteur, 1964  
*bernardi* FitzSimons, 1958 (*bonisi*\* Pasteur, 1962)  
*blanci* Pasteur, 1967  
*bradfieldi* Hewitt, 1932  
*capensis* A. Smith, 1849 (*strigatus* Gray, 1864;  
*grotei*\* Sternfeld, 1911; *ngamiensis*  
 FitzSimons, 1932; *pakenhami*\* Loveridge, 1941)  
*chobiensis* FitzSimons, 1932  
*conradti* Matschie, 1892  
*conraui* Tornier, 1902 (*strongi* Barbour and  
 Loveridge, 1927)  
*decaryi* Angel, 1930  
*depressus* K. P. Schmidt, 1919  
*expectatus* Pasteur and Blanc, 1967  
*fischeri* Boulenger, 1890  
*grandisonae* Pasteur, 1962  
*gravis* Pasteur, 1964  
*guibei* Pasteur, 1964  
*gutturalis* Barboza du Bocage, 1873 (*paurospilus*\*  
 Laurent, 1952; *dysmicus*\* Perret, 1963)  
*heterurus* Boettger, 1913  
*inexpectatus* Pasteur, 1964  
*insularis* Boettger, 1913  
*keniensis* Parker, 1936  
*klemmeri* Pasteur, 1964  
*klugei* H. M. Smith, Martin and Swain, 1977  
*lawrencei* Hewitt, 1926  
*luteopicturatus* Pasteur, 1964 (*zanzibaritis*\*)

- Pasteur, 1964)  
*madagascariensis* Boettger, 1881 (*pictus* W. Peters, 1883; *petteri*\* Pasteur and Blanc, 1967)  
*manni* Loveridge, 1928  
*methueni* FitzSimons, 1937  
*miops* Günther, 1891 (*spinulifer* Boettger, 1913)  
*montanus* Pasteur, 1964  
*ocellatus* Roux, 1907  
*ornatus* Pasteur, 1964  
*picturatus* W. Peters, 1868 (*variegatus* W. Peters, 1868; *griseus* Toriner, 1896; *septemlineatus* Toriner, 1896; *quinquelineatus* Tornier, 1896; *mombasicus*\* Loveridge, 1935; *sudanensis*\* Loveridge, 1935; *ukerewensis* Loveridge, 1935)  
*pictus* W. Peters, 1883 (*robustus* Boettger, 1913)  
*rarus* Pasteur and Blanc, 1973  
*rex* Broadly, 1963  
*scheffleri* Sternfeld, 1912 (*laterimaculatus*\* Pasteur, 1964; *compositus*\* Pasteur, 1964; *ulugurensis*\* Pasteur, 1964)  
*scorteccii* Pasteur, 1959  
*septemtuberculatus* Angel, 1942  
*somalicus* Loveridge, 1931 (*fisheri* Scortecci, 1931; *annectans* Loveridge, 1935; *battersbyi*\* Pasteur, 1962; *howelli*\* Pasteur and Broadly, 1988)  
*stevensoni* Hewitt, 1926  
*thomensis* W. Peters, 1880 (*rolasi* Greef, 1885; *delicatus*\* Pasteur, 1962; *wermuthi*\* Pasteur, 1962)  
*tolampyae* Grandidier, 1872 (*tuberifer* Boettger, 1913)  
*tuberosus* Mertens, 1965  
*verticillatus* Mocquard, 1895  
*wetzeli* H. M. Smith, Martin and Swain, 1977  
*williamsi* Loveridge, 1952
- Mediodactylus* Shcherbak and Golubev, 1977  
*amictopholis* Hoofien, 1967  
*brevipes* Blanford, 1874  
*heterocercus* Blanford, 1874 (*mardinensis*\* Mertens, 1924)  
*kotschyi* Steindachner, 1870 (*cyprius* Fitzinger, 1843; *concolor*\* Bedriaga, 1882; *maculatus*\* Bedriaga, 1882; *danilewskii*\* Strauch, 1887; *oertzeni*\* Boettger, 1888; *colchicus*\* Nikolsky, 1902; *bartoni*\* Stepanek, 1934; *fitzingeri*\* Stepanek, 1937; *orientalis*\* Stepanek, 1937; *solerii*\* Wettstein, 1937; *steindachneri*\* Stepanek, 1937; *stepaneki*\* Wettstein, 1937; *syriacus*\* Stepanek, 1937; *unicolor* Wettstein, 1937; *wettsteini*\*

Stepanek, 1937; *buresschi* Stepanek, 1937;  
*rumelicus*\* L. Müller, 1939; *saronicus*\*  
 Werner, 1939; *kalypsae*\* Stepanek, 1939;  
*lycaonicus*\* Mertens, 1952; *rarus* Wettstein,  
 1952; *stubbei* Wettstein, 1952; *christianae*  
 Buchholz, 1955; *skopjensis*\* Karaman, 1965;  
*buchholzi*\* Beutler and Gruber, 1977;  
*schultzewestrumi*\* Beutler and Gruber, 1977;  
*gruberi* Beutler and Gruber, 1977; *fuchsi*\*  
 Beutler and Gruber, 1977; *bibroni*\* Beutler  
 and Gruber, 1977; *adelphiensis*\* Beutler and  
 Gruber, 1978; *bileki* Tiedemann and Haupl,  
 1980; *tinensis*\* Beutler and Fror, 1980;  
*beutleri*\* Baran and Gruber, 1981; *karabagi*\*  
 Baran and Gruber, 1982; *ciliciensis*\* Baran  
 and Gruber, 1982; *ponticus*\* Baran and Gruber,  
 1982)

*russowii* Strauch, 1887 (*zarudnyi*\* Nikolsky, 1899;  
*copalensis* Schnitnikow, 1928)  
*sagittifer* Nikolsky, 1899  
*spinicauda* Nikolsky, 1887

*Microgecko* Nikolsky, 1907

*helenae* Nikolsky, 1907 (*fasciatus*\* Schmidtler and  
 Schmidtler, 1972)  
*latifi* Leviton and S. Anderson, 1972  
*persicus* Nikolsky, 1903 (*bakhtiari*\* Minton, S.  
 Anderson, J. A. Anderson, 1970;  
*euphorbiacola*\* Minton, S. Anderson, and J. A.  
 Anderson, 1970)

*Microscalabotes* Boulenger, 1883

*bivittis* W. Peters, 1883 (*hildebrandti* W. Peters,  
 1883; *cowanii* Boulenger, 1883)

*Millotisaurus* Pasteur, 1962

*mirabilis* Pasteur, 1962

*Nactus* Kluge, 1983 (*Mascarenogecko* Ulber and Gericke, 1988)

*pelagicus* Girard, 1858 (*arnouxii* A. Duméril, 1851;  
*multicarinatus* Günther, 1872; *arfakianus*  
 Meyer, 1874; *fasciata* Macleay, 1877;  
*marmorata* Macleay, 1877; *eboracensis* Macleay,  
 1877; *cheverti* Boulenger, 1885; *heteronotus*  
 Boulenger, 1885; *undulatus*\* Kopstein, 1926)  
*coindermirensis* Bullock, Arnold and Bloxan, 1985  
*galgajuga* Ingram, 1978  
*serpensinsula* Loveridge, 1951  
*vankampeni* Brongersma, 1933

*Narudasia* Methuen and Hewitt, 1914

*festiva* Methuen and Hewitt, 1914

- Naultinus* Gray, 1842 (*Heteropholis* Fischer, 1882;  
*Naultinulus* Chrapliwy, H. M. Smith, and  
 Grant, 1961)  
*elegans* Gray, 1842 (*punctatus\** Gray, 1843;  
*lineatus* Gray, 1869; *sulphurus* Buller, 1871;  
*pentagonalis* Colenso, 1880; *ocellatus* McCann,  
 1955)  
*gemmeus* McCann, 1955  
*grayii* Bell, 1843 (*simpsoni* Robb, 1980)  
*manukanus* McCann, 1955  
*poecilochlorus* Robb, 1980  
*rudis* Fischer, 1881 [1882]  
*stellatus* Hutton, 1872 (*pulcherrimus* Buller, 1877)  
*tuberculatus* McCann, 1955
- Nephrurus* Günther 1876 (*Underwoodisaurus* Wermuth, 1965)  
*asper* Günther, 1876  
*deleani* Harvey, 1983  
*laevissimus* Mertens, 1958  
*laevis* De Vis, 1886 (*platyurus* Boulenger, 1886;  
*occidentalis\** Storr, 1963; *pilbarensis\**  
 Storr, 1963)  
*milii* Bory de Saint-Vincent, 1825 (*miliusii*  
 Duméril and Bibron, 1836; *blavieri* de  
 Rochebrune, 1884; *asper* Boulenger, 1913;  
*husbandi* Wells and Wellington, 1984)  
*sphyrurus* Ogilby, 1892 (*walshi* Kinghorn, 1931)  
*stellatus* Storr, 1968  
*vertebralis* Storr, 1963  
*wheeleri* Loveridge, 1932 (*cinctus\** Storr, 1963)
- Oedura* Gray, 1842 (*Pachyurus* Fitzinger, 1843; *Amalosis*  
 Wells and Wellington, 1984)  
*castelnaui* Thomillot, 1889 (*mayeri* Garman, 1901)  
*coggeri* Bustard, 1966  
*filicipoda* King, 1984  
*gemmata* King and Gow, 1983  
*gracilis* King, 1984  
*lesueurii* Duméril and Bibron, 1836 (*phillipsi*  
 Wells and Wellington, 1985)  
*marmorata* Gray, 1842 (*fracticolor* De Vis, 1884;  
*cincta* De Vis, 1888; *attenboroughi* Wells and  
 Wellington, 1985; *derelicta* Wells and  
 Wellington, 1985; *greeri* [n.n.] Wells and  
 Wellington, 1985)  
*monilis* De Vis, 1888  
*obscura* King, 1984  
*reticulata* Bustard, 1969  
*rhombifer* Gray, 1845  
*robusta* Boulenger, 1885  
*tryoni* De Vis, 1884 (*ocellata* Boulenger, 1885)

*verrilli* [?] Cope, 1868

*Ophidiocephalus* Lucas and Frost, 1897

*taeniatus* Lucas and Frost, 1897

*Pachydactylus* Wiegmann, 1834 (*Colobopus* Fitzinger, 1843; *Cantina* Gray, 1845; *Homodactylus* Gray, 1864; *Elasmodactylus* Boulenger, 1894)

*amoenus* Werner, 1910

*austeni* Hewitt, 1923

*bibronii* A. Smith, 1845 [1846] (*turneri* Gray, 1864; *stellatus* Werner, 1910; *pulitzerae*\* K. P. Schmidt, 1933)

*bicolor* Hewitt, 1926

*capensis* A. Smith, 1846 (*elegans* Gray, 1845; *mentomarginatus* A. Smith, 1849; *oshaughnessyi*\* Boulenger, 1885; *obscurus* Thomiot, 1889; *affinis*\* Boulenger, 1896; *tessellatus* Werner, 1910; *leopardinus* Sternfeld, 1911; *levyi* FitzSimons, 1933; *vansoni*\* FitzSimons, 1933; *katanganus*\* DeWitte, 1953)

*caraculicus* FitzSimons, 1959

*fasciatus* Boulenger, 1888

*formosus* [?] A. Smith, 1849

*gaiasensis* Steyn and Mitchell, 1967

*geitje* Sparrman, 1778 (*ocellatus* Cuvier, 1817; *inunguis* Cuvier, 1817; *bergii* Wiegmann, 1834)

*kobosensis* FitzSimons, 1938

*kochii* FitzSimons, 1959

*labialis* FitzSimons, 1938

*laevigatus* Fischer, 1888 (*tessellatus* FitzSimons, 1938; *fitzsimonsi*\* Loveridge, 1947)

*maculatus* Gray, 1845 (*albomarginatus* Hewitt, 1932; *microlepis* Hewitt, 1935)

*mariquensis* A. Smith, 1849 (*latirostris*\* Hewitt, 1923; *macrolepis* FitzSimons, 1939)

*mentalis* Hewitt, 1926

*monticolus* [?] FitzSimons, 1943

*namaquensis* Sclater, 1898

*oculatus* Hewitt, 1927

*oreophilus* McLachlan and Spence, 1967

*punctatus* W. Peters, 1854 (*brunnthaleri* Werner, 1913; *langi* FitzSimons, 1932; *amoenoides*\* Hewitt, 1935; *scherzi*\* Mertens, 1954)

*rugosus* A. Smith, 1849 (*frater* Hewitt, 1935; *barnardi*\* FitzSimons, 1941)

*sansteyni* Steyn and Mitchell, 1967

*scutatus* Hewitt, 1927 (*robertsi*\* FitzSimons, 1938; *angolensis*\* Loveridge, 1944)

*serval* Werner, 1910 (*purcelli*\* Boulenger, 1910;

*pardus* Sternfeld, 1911; *montanus* Methuen and



- Hewitt, 1914; *onscepensis*\* Hewitt, 1935)  
*tetensis* Loveridge, 1953  
*tigrinus* Van Dam, 1921 (*rhodesianus* Loveridge,  
 1947)  
*tsodiloensis* Haacke, 1966  
*tuberculosis* Boulenger, 1894 (*triedrus* Boulenger,  
 1913; *boulengeri* Toriner, 1896)  
*weberi* Roux, 1907 (*gariesensis* Hewitt, 1932;  
*wernerii*\* Hewitt, 1935; *acuminatus*\*  
 FitzSimons, 1941)
- Palmatogeocko* Andersson, 1908 (*Syndactylosaura* Werner, 1910)  
*rangei* Andersson, 1908 (*schultzei* Werner, 1910)
- Paradelma* Kinghorn, 1926  
*orientalis* Günther, 1876
- Paragehyra* Angel, 1929  
*petiti* Angel, 1929
- Paroedura* Günther, 1879  
*androyensis* Grandidier, 1867 (*porogaster*  
 Boulenger, 1896)  
*bastardi* Mocquard, 1900  
*gracilis* Boulenger, 1896  
*guibeae* Dixon and Kroll, 1974  
*homalorhinus* Angel, 1936  
*oviceps* Boettger, 1881  
*pictus* W. Peters, 1854 (*madagascariensis* Mocquard,  
 1895; *robustus* Boulenger, 1896)  
*sanctijohannis* Günther, 1879  
*stumpffi* Boettger, 1879
- Perochirus* Boulenger, 1885  
*ateles* A. Duméril, 1856 (*articulatus* Fischer,  
 1882; *depressus* Fischer, 1882; *klugei*  
 Wermuth, 1965)  
*guentheri* Boulenger, 1885  
*scutellatus* Fischer, 1882
- Phelsuma* Gray, 1825 (*Anoplopus* Wagler, 1830; *Chiroperus*  
 Wiegmann, 1834; *Rhoptropella* Hewitt, 1937;  
*Macrophelsuma* Hoffstetter, 1946;  
*Archaeophelsuma* Borner, 1972; *Neophelsuma*  
 Borner, 1972)  
*abbotti* Stejneger, 1893 (*sumptio*\* Cheke, 1982)  
*andamanensis* Blyth, 1860 (*chamaelon* Tytler, 1864)  
*astriata* Tornier, 1901 (*carinatum* [?] Rendahl,  
 1939; *maheense* Rendahl, 1939; *astovei*\*  
 FitzSimons, 1948; *semicarinata*\* Cheke, 1982)  
*barbouri* Loveridge, 1942  
*befotakensis* Borner and Minuth, 1982

- borbonica* Mertens, 1966 (*agalegae*\* Cheke, 1975)  
*cepediana* Merrem, 1820  
*chekei* Borner and Minuth, 1982  
*comorensis* Boettger, 1913  
*dubia* Boettger, 1881  
*flavigularis* Mertens, 1962  
*gigas* Lienard, 1842 (*newtonii* Günther, 1877;  
*newtonii* Boulenger, 1884; *edwardnewtonii*  
Vinson and Vinson, 1969)  
*guentheri* Boulenger, 1885  
*guimbeaui* Mertens, 1963 (*rosagularis*\* Vinson and  
Vinson, 1969)  
*guttata* Kaudern, 1922  
*laticauda* Boettger, 1880 (*angularus*\* Mertens,  
1964)  
*lineata* Gray, 1842 (*bifasciata*\* Boettger, 1913;  
*chloroscelis*\* Mertens, 1962; *bombetokensis*\*  
Mertens, 1964; *dorsivittata*\* Mertens, 1964;  
*pusilla*\* Mertens, 1964; *punctulata*\* Mertens,  
1970; *leiogaster* Mertens, 1973)  
*madagascariensis* Gray, 1831 (*tetradactylus* [?]  
Schneider, 1797; *sarroubea* [?] Daudin, 1802;  
*grandis*\* Gray, 1870; *kochi*\* Mertens, 1954;  
*martensi* Mertens, 1962; *venusta*\* Mertens,  
1964; *notissima* Mertens, 1970; *boehmei*\*  
Meier, 1982)  
*minuthi* Borner, 1980  
*modesta* Mertens, 1970  
*mutabilis* Grandidier, 1869 (*breviceps*\* Boettger,  
1894; *androyense* Mocquard, 1901; *micropholis*  
Boettger, 1913)  
*nigristriata* Meier, 1984  
*ocellata* Boulenger, 1885  
*ornata* Gray, 1825 (*vinsoni* Mertens, 1963;  
*inexpectata*\* Mertens, 1966)  
*parkeri* Loveridge, 1941  
*quadriocellata* W. Peters, 1883 (*bimaculata*\*  
Kaudern, 1922; *leiura*\* Meier, 1983; *parva*\*  
Meier, 1983)  
*robertmertensi* Meier, 1980  
*serraticauda* Mertens, 1963  
*standingi* Methuen and Hewitt, 1913  
*sundbergi* Rendahl, 1939 (*longinsulae*\* Rendahl,  
1939; *pulchra* Rendahl, 1939; *cousinense*  
Rendahl 1939; *menaiensis* Mertens, 1966;  
*ladiguensis*\* Bohme and Meier, 1981; *rubra*  
Borner and Minuth, 1982; *umbrae* Borner and  
Minuth, 1982)  
*trilineata* Gray, 1842  
*v-nigra* Boettger, 1913 (*pasteuri*\* Meier, 1984)

- Phyllodactylus* Gray, 1828 (*Euleptes* Fitzinger, 1843;  
*Discodactylus* Fitzinger, 1843; *Gerrhopygus*  
 Gray, 1845)  
*angelensis* Dixon, 1966  
*angustidigitus* Dixon and Huey, 1970  
*ansorgii* Boulenger, 1907  
*apricus* Dixon, 1966  
*barringtonensis* Van Denburgh, 1912  
*baurii* Garman, 1892 (*gorii*\* Lanza, 1973)  
*bordai* Taylor, 1942  
*bugastrolepis* Dixon, 1966  
*clinatus* Dixon and Huey, 1970  
*darwini* Taylor, 1942  
*davisi* Dixon, 1964  
*delcampi* Mosauer, 1936  
*dixoni* Rivero-Blanco and Lancini, 1968  
*duellmani* Dixon, 1960  
*europaeus* Gene, 1839 (*wagleri* Fitzinger, 1843;  
*doriae* Lataste, 1877)  
*galapagensis* W. Peters, 1869 (*daphnensis*\* Van  
 Denburgh, 1912; *duncanensis* Van Denburgh,  
 1912; *maresi*\* Lanza, 1973; *olschkii*\* Lanza,  
 1973)  
*gerrhopygus* Wiegmann, 1835 (*gymnopygus* Duméril and  
 Bibron, 1836)  
*gilberti* Heller, 1903  
*heterurus* Werner, 1907  
*homolepidurus* H. M. Smith, 1935 (*nolascoensis*\*  
 Dixon, 1964)  
*inaequalis* Cope, 1875  
*insularis* Dixon, 1960  
*interandinus* Dixon and Huey, 1970  
*johnwrighti* Dixon and Huey, 1970  
*julieni* Cope, 1885  
*kofordi* Dixon and Huey, 1970  
*lanei* H. M. Smith, 1935 (*magnatuberculatus* Taylor,  
 1940; *rupinus*\* Dixon, 1964)  
*leei* Cope, 1889  
*lepidopygus* Tschudi, 1845 (*phacophorus* Tschudi,  
 1845; *nigrofasciatus* Cope, 1878; *variegatus*  
 Werner, 1900)  
*lineatus* Gray, 1838 (*essexi*\* Hewitt, 1925;  
*rupicolus*\* FitzSimons, 1938)  
*martini* Lidth de Jeude, 1887  
*melanostictus* Taylor, 1962  
*microlepidotus* FitzSimons, 1939  
*microphyllus* Cope, 1876  
*muralis* Taylor, 1939 (*isthmus*\* Dixon, 1964)  
*palmeus* Dixon, 1968  
*partidus* Dixon, 1966  
*paucituberculatus* Dixon, 1960

- peringueyi* [?] Boulenger, 1910  
*porphyreus* Daudin, 1802 (*brevipes* Mocquard, 1900;  
*namaquensis* Hewitt, 1935; *cronwrighti* Hewitt,  
 1937)  
*pulcher* Gray, 1828 (*spatulatus* Cope, 1863)  
*pumilis* Dixon and Huey, 1970  
*reissii* W. Peters, 1862 (*baessleri* Werner, 1900;  
*guayaquilensis* Werner, 1910; *abrupteseriatus*  
 Werner, 1913; *magister* Noble, 1924)  
*riebeckii* W. Peters, 1882  
*rutteni* Hummelinck, 1947  
*santacruzensis* Dixon, 1966  
*sentosus* Dixon and Huey, 1970  
*siamensis* Boulenger, 1898 (*paviei* Mocquard, 1904;  
*burmanicus* Annandale, 1905)  
*tinklei* Dixon, 1966  
*trachyrhinus* Boulenger, 1899  
*transversalis* Huey, 1975  
*tuberculosis* Wiegmann, 1835 (*magnus\** Taylor, 1940;  
*edwardofischeri* Mertens, 1952; *ingeri\** Dixon,  
 1964; *saxatilis\** Dixon, 1964)  
*unctus* Cope, 1864  
*ventralis* O'Shaughnessy, 1875 (*mulleri* Parker,  
 1935; *underwoodi* Dixon, 1962)  
*wirshingi* Kerster and H. M. Smith, 1955  
 (*hispaniolae\** Schwartz, 1979; *sommeri\**  
 Schwartz, 1979)  
*xanti* Cope, 1864 (*mentalis* Werner, 1910;  
*nocticolus\** Dixon, 1964; *zweifeli\** Dixon,  
 1964; *acorius\** Dixon, 1966; *angulus\** Dixon,  
 1966; *circus\** Dixon, 1966; *estabanensis\**  
 Dixon, 1966; *coronatus\** Dixon, 1966; *sloani*  
 Bostic, 1971)
- Phyllopezus* W. Peters, 1877  
*periosus* Rodrigues, 1986  
*pollicaris* Spix, 1825 (*goyazensis* W. Peters, 1877;  
*przewalskii\** Koslowsky, 1894; *spixii*  
 Schlegel, 1933)
- Phyllurus* Goldfuss, 1820 (*Geckoides* Peron, 1807;  
*Phyllurus* [n.n.] Oken, 1817; *Phyllura* Kaup,  
 1827; *Anomalurus* Fitzinger, 1843)  
*caudiannulatus* Covacevich, 1975  
*cornutus* Ogilby, 1892 (*lichenosus* Günther, 1897;  
*swaini* Wells and Wellington, 1985)  
*platurus* Shaw, 1790 (*phyllurus* Schneider, 1797;  
*discosura* Merrem, 1820; *platyura* Merrem,  
 1820; *novae-hollandiae* Cuvier, 1822; *whitii*  
 Gray, 1825; *cuvieri* Bory de Saint-Vincent,  
 1828; *platycaudus* Schinz, 1835; *australis*  
 Swainson, 1839; *inermis* Gray, 1845)

*salebrosus* Covacevich, 1975

*Pletholax* Cope, 1864

*gracilis* Cope, 1864 (*edeleusis*\* Storr, 1978)

*Pristurus* Rüppell, 1835 (*Spatalura* Gray, 1863;

*Geisoprysturus* Lanza and Sassi, 1968)

*abdelkuri* Arnold, 1986

*carteri* Gray, 1863 (*tuberculatus*\* Parker, 1931)

*celerrimus* Arnold, 1977

*collaris* Steindachner, 1867

*crucifer* Valenciennes, 1861 (*longipes* W. Peters,

1871; *stefaninii* Calabresi, 1927;

*laticephalus* Scortecci, 1933)

*flavipunctatus* Rüppell, 1835 (*percristatus*

Boulenger, 1896; *pseudoflavipunctatus*

Scortecci, 1933)

*gasperetti* Arnold, 1986 (*gallagheri*\* Arnold, 1986)

*guichardi* Arnold, 1986

*insignis* Blanford, 1881

*insignoides* Arnold, 1986

*minimus* Arnold, 1977

*ornithocephalus* Arnold, 1986

*phillipsii* Boulenger, 1895 (*somalicus* Parker,

1932)

*popovi* Arnold, 1982

*rupestris* Blanford, 1874 (*migiurtinicus* Scortecci,

1933; *guweirensis*\* Haas, 1951; *iranicus*\* K.

P. Schmidt, 1952)

*saada* Arnold, 1986

*simonettai* Lanza and Sassi, 1968

*sokotranus* Parker, 1938

*Pseudogekko* Taylor, 1922

*brevipes* Boettger, 1897

*compressicarpus* Taylor, 1915

*labialis* W. Peters, 1867

*smaragdinus* Taylor, 1922

*Pseudogonatodes* Ruthven, 1915

*barbouri* Noble, 1921

*furvus* Ruthven, 1915

*guianensis* Parker, 1935 (*amazonicus*\* Vanzolini,

1967)

*lunulatus* Roux, 1927

*peruvianus* Huey and Dixon, 1970

*Ptenopus* Gray, 1865

*carpi* Brain, 1962

*garrulus* A. Smith, 1849 (*maculatus*\* Gray, 1865)

*kochi* Haacke, 1964

- Ptychozoon* Kuhl, 1822 (*Ptyxozoon* Kuhl and Hasselt, 1822; *Pteropleura* Gray, 1827)  
*horsfieldii* Gray, 1827  
*intermedium* Taylor, 1915  
*kuhli* Stejneger, 1902 (*homalocephala* Creveldt, 1809)  
*lionotum* Annandale, 1905  
*rhacophorus* Boulenger, 1899
- Ptyodactylus* Goldfuss, 1820  
*guttatus* Heyden, 1827 (*maculatus* Schinz, 1833; *lacazzi* Boutan, 1892; *siphonorhina* J. Anderson, 1896)  
*hasselquistii* Donndorff, 1798 (*ascalabotes* Merrem, 1820; *lobatus* Geoffroy, 1823; *ragazzi*\* J. Anderson, 1898; *togoensis* Tornier, 1901)  
*homolepis* Blanford, 1876 (*socotranus*\* Steindachner, 1902)  
*oudrii* Lataste, 1880  
*puiseuxi* Bouton, 1893 (*bischoffsheimi* Boutan, 1893; *montmahoui* Bouton, 1893; *barroisi* Bouton, 1893; *syriacus* Peracca, 1894; *sancti-montis* Barbour, 1914)
- Pygopus* Merrem, 1820 (*Hysteropus* A. Duméril, in Duméril and Bibron, 1839; *Cryptodelma* Fischer, 1882)  
*lepidopodus* Lacepede, 1804 (*novaehollandiae* Oppel, 1811; *lepidopus* Merrem, 1820; *squamiceps* Gray, 1845; *longicaudatus* Tepper, 1882)  
*nigriceps* Fischer, 1882 (*baileyi* Günther, 1897; *schraderi*\* Boulenger, 1913; *klugei* Wells and Wellington, 1985; *territorianus* [n.n.] Wells and Wellington, 1985)
- Quedenfeldtia* Boettger, 1883  
*trachyblepharus* Boettger, 1874 (*moerens* Chabanaud, 1916)
- Rhacodactylus* Fitzinger, 1843 (*Correlophus* Guichenot, 1866; *Ceratolophus* Barboza du Bocage, 1873; *Chamaeleonurus* Boulenger, 1878; *Chamaeleonurus* Carus, 1889; *Pseudothecadactylus* Brongersma, 1936; *Torresia* Brongersma, 1934)  
*auriculatus* Bavay, 1869 (*hexaceros* Barboza du Bocage, 1873)  
*australis* Günther, 1877  
*cavaticus* Cogger, 1975  
*chahoua* Bavay, 1869  
*ciliatus* Guichenot, 1866  
*leachianus* Cuvier, 1829 (*aubryanus* Barboza du Bocage, 1873)  
*lindneri* Cogger, 1975

*sarasinorum* Roux, 1913  
*trachyrhynchus* Barboza du Bocage, 1873  
 (*trachycephalus* Boulenger, 1878)

*Rhoptropus* W. Peters, 1869 (*Dactychylikion* Thominot, 1878)  
*afer* W. Peters, 1869  
*barnardi* Hewitt, 1926  
*biporosus* FitzSimons, 1957  
*boultoni* K. P. Schmidt, 1933 (*benguellensis*\*  
 Mertens, 1938; *montanus*\* Laurent, 1964)  
*braconnieri* Thominot, 1878  
*bradfieldi* Hewitt, 1935 (*diporus*\* Haacke, 1965)  
*taeniostictus* Laurent, 1964

*Rhynchoedura* Günther, 1867  
*ornata* Günther, 1867 (*paraornatus* Wells and  
 Wellington, 1984; *ormsbyi* Wells and  
 Wellington, 1985)

*Saurodactylus* Fitzinger, 1843  
*fasciatus* Werner, 1931  
*mauritanicus* Duméril and Bibron, 1836 (*desertorum*  
 Fitzinger, 1843; *brosseti*\* Bons and Pasteur,  
 1957)

*Sphaerodactylus* Wagler, 1830 (*Sphaeriodactylus* Gray, 1831)  
*alayoi* Grant, 1959  
*altavelensis* Noble and Hassler, 1933  
 (*brevirostratus*\* Shreve, 1968;  
*enrigoilloensis*\* Shreve, 1968; *lucioi* Thomas  
 and Schwartz, 1983)  
*argivus* Garman, 1888 (*bartschi*\* Cochran, 1934;  
*lewisi*\* Grant, 1941)  
*argus* Gosse, 1850 (*henriquesi* Grant, 1940;  
*andresensis*\* Dunn and Saxe, 1950)  
*armasi* Schwartz and Garrido, 1974  
*armstrongi* Noble and Hassler, 1933 (*hypsinephes*\*  
 Thomas and Schwartz, 1983)  
*asterulus* Schwartz and Graham, 1980  
*beattyi* Grant, 1937 (*seamani*\* Thomas and Schwartz,  
 1966)  
*becki* K. P. Schmidt, 1919  
*bromeliarum* G. Peters and Schwartz, 1972  
*caicosensis* Cochran, 1934  
*callocricus* Schwartz, 1976  
*cinereus* Wagler, 1830  
*clenchi* Shreve, 1968 (*apocoptus*\* Schwartz, 1983)  
*cochranae* Ruibal, 1946  
*copei* Steindachner, 1867 (*anthracinus* Cope, 1862;  
*picturatus*\* Garman, 1887; *asper* Garman, 1888;  
*cataplexis*\* Schwartz and Thomas, 1965;  
*enochrus*\* Schwartz and Thomas, 1965;

- polyommatus*\* Thomas, 1968; *astreptus*\*  
 Schwartz, 1975; *pelates*\* Schwartz, 1975;  
*websteri*\* Schwartz, 1975; *deuterus*\* Schwartz,  
 1975)  
*corticola* Garman, 1888 (*aporrox*\* Schwartz, 1968;  
*campter*\* Schwartz, 1968; *soter*\* Schwartz,  
 1968)  
*cryphius* Thomas and Schwartz, 1977  
*darlingtoni* Shreve, 1968 (*noblei*\* Shreve, 1968;  
*bobilini*\* Thomas and Schwartz, 1983;  
*mekistus*\* Thomas and Schwartz, 1983)  
*difficilis* Barbour, 1914 (*lycauges*\* Schwartz,  
 1983; *euopter*\* Schwartz, 1983; *typhlops*\*  
 Schwartz, 1983; *peratus*\* Schwartz, 1983;  
*diolenius*\* Schwartz, 1983; *anthracomus*\*  
 Schwartz, 1983)  
*dunni* K. P. Schmidt, 1936  
*elasmorhynchus* Thomas, 1966  
*elegans* Macleay, 1834 (*punctatissimus* Duméril and  
 Bibrion, 1836; *alopex* Cope, 1862)  
*elegantulus* Barbour, 1917  
*fantasticus* Duméril and Bibrion, 1836  
 (*ligniservulus*\* King, 1962; *anidrotus*\*  
 Thomas, 1965; *fuga*\* Thomas, 1965; *hippomanes*\*  
 Thomas, 1965; *karukera*\* Thomas, 1965;  
*orescius*\* Thomas, 1965; *phyzacinus*\* Thomas,  
 1965; *tartaropylorus*\* Thomas, 1965)  
*gaigeae* Grant, 1932  
*gilvitorques* Cope, 1862  
*glaucus* Cope, 1865 (*inornatus* W. Peters, 1873;  
*torquatus* Strauch, 1887)  
*goniorhynchus* Cope, 1895  
*graptolaemus* Harris and Kluge, 1983  
*heliconiae* Harris, 1982  
*homolepis* Cope, 1886 (*carinatus* Andersson, 1916;  
*imbricatus* Andersson, 1916; *mertensi* Wermuth,  
 1965)  
*inaguae* Noble and Klingel, 1932  
*intermedius* Barbour and Ramsden, 1919 (*drapetiscus*  
 Schwartz, 1958)  
*klauberi* Grant, 1931  
*ladae* Thomas and Hedges, 1988  
*lazelli* Shreve, 1968  
*leucaster* Schwartz, 1973  
*levinsi* Heatwole, 1968  
*lineolatus* Lichtenstein and von Martens, 1856  
 (*casicolus* Cope, 1862)  
*macrolepis* Günther, 1859 (*grandisquamis* Stejneger,  
 1904; *danforthi* Grant, 1931; *parvus*\* King,  
 1962; *ateles*\* Thomas and Schwartz, 1966;  
*guarionex*\* Thomas and Schwartz, 1966; *inigoii*\*  
 Thomas and Schwartz, 1966; *mimetes*\* Thomas



and Schwartz, 1966; *phoberus*\* Thomas and Schwartz, 1966; *spanius*\* Thomas and Schwartz, 1966; *stibarus*\* Thomas and Schwartz, 1966)  
*mariguanae* Cochran, 1934  
*microlepis* Reinhardt and Lütken, 1863  
 (*melanospilos* Bocourt, 1873; *thomasi*\* Schwartz, 1965)  
*micropithecus* Schwartz, 1977  
*millepunctatus* Hallowell, 1861 (*continentalis* Werner, 1896)  
*molei* Boettger, 1894 (*buergeri* Werner, 1900; *venezuelanus* Roux, 1927; *boettgeri* [n.n.] Donoso Barros, 1968)  
*monensis* Meerwarth, 1901  
*nicholsi* Grant, 1931  
*nigropunctatus* Gray, 1845 (*decoratus*\* Garman, 1888; *flavicauda*\* Barbour, 1904; *gibbus*\* Barbour, 1921; *atessares* Thomas and Schwartz, 1966; *granti*\* Thomas and Schwartz, 1966; *lissodesmus*\* Thomas and Schwartz, 1966; *strategus*\* Thomas and Schwartz, 1966; *porrasi*\* Schwartz, 1972)  
*notatus* Baird, 1858 (*exsul*\* Barbour, 1914; *amaurus*\* Schwartz, 1966; *atactus*\* Schwartz, 1966; *peltastes*\* Schwartz, 1966)  
*nycteropus* Thomas and Schwartz, 1977  
*ocoae* Schwartz and Thomas, 1977  
*oliveri* Grant, 1944 (*storeyae*\* Grant, 1944)  
*omoglaux* Thomas, 1982  
*oxyrhinus* Gosse, 1850 (*dacnicolor*\* Barbour, 1910)  
*pacificus* Stejneger, 1903  
*parkeri* Grant, 1939  
*parthenopion* Thomas, 1965  
*perissodactylius* Thomas and Hedges, 1988  
*ramsdeni* Ruibal, 1959  
*randi* Shreve, 1968 (*methorius*\* Schwartz, 1977; *strahmi*\* Schwartz, 1977)  
*rhabdotus* Schwartz, 1970  
*richardsoni* Gray, 1845 (*gossei*\* Grant, 1939)  
*roosevelti* Grant, 1931  
*rosaurae* Parker, 1940  
*ruibali* Grant, 1959  
*sabanus* Cochran, 1938  
*samanensis* Cochran, 1932  
*savagei* Shreve, 1968 (*juanilloensis*\* Shreve, 1968)  
*scaber* Barbour and Ramsden, 1919  
*scapularis* Boulenger, 1902  
*semasiops* Thomas, 1975  
*shrevei* Lazell, 1961  
*sommeri* Graham, 1981  
*sputator* Sparrman, 1784 (*pictus* Garman, 1888)  
*streptophorus* Thomas and Schwartz, 1977

(*sphenophanes*\* Thomas and Schwartz, 1983)  
*stejnegeri* Cochran, 1931  
*thompsoni* Schwartz and Franz, 1976  
*torrei* Barbour, 1914 (*spielmani*\* Grant, 1958;  
*ocujal*\* Thomas and Schwartz, 1966)  
*townsendi* Grant, 1931  
*underwoodi* Schwartz, 1968  
*vincenti* Boulenger, 1891 (*festus*\* Barbour, 1915;  
*monilifer*\* Barbour, 1921; *adamas*\* Schwartz,  
 1965; *diamesus*\* Schwartz, 1965; *josephinae*\*  
 Schwartz, 1965; *pheristus*\* Schwartz, 1965;  
*psammius*\* Schwartz, 1965; *ronaldi*\* Schwartz,  
 1965)  
*williamsi* Thomas and Schwartz, 1983  
*zygaena* Schwartz and Thomas, 1977

*Stenodactylus* Fitzinger, 1826 (*Goniodactylus* Schlegel,  
 1826; *Tolarenta* Gray, 1842; *Ceramodactylus*  
 Blanford, 1872; *Pseudoceramodactylus* Haas,  
 1957; *Trigonodactylus* Haas, 1957; *Garzoniella*  
 Perret, 1976)  
*affinis* Murray, 1884  
*arabicus* Haas, 1957  
*doriae* Blanford, 1874 (*major* Parker, 1930)  
*grandiceps* Haas, 1952  
*khobarensis* Haas, 1957  
*leptocosymbotes* Leviton and S. Anderson, 1967  
*petrii* J. Anderson, 1896 (*stenurus* Werner, 1899;  
*elimensis* Barbour, 1914)  
*pulcher* J. Anderson, 1896  
*slevini* Haas, 1957 (*arabicus* Haas, 1957; *hassi*  
 Kluge, 1967)  
*sthenodactylus* Lichtenstein, 1823 (*elegans*  
 Fitzinger, 1826; *savignyi* Audouin, 1827;  
*guttatus* Cuvier, 1829; *wilkinsonii* Gray,  
 1842; *mauritanicus* Guichenot, 1850; *hirouxii*  
 Doumergue, 1899; *longipes* Perret, 1976)  
*yemenensis* Arnold, 1980

*Strophurus* Fitzinger, 1843 (*Strophura* Gray, 1845;  
*Oedurella* Lönnberg and Andersson, 1913;  
*Eremiastrophurus* Wells and Wellington, 1985)  
*assimilis* Storr, 1988  
*ciliaris* Boulenger, 1885 (*aberrans*\* Glauert, 1952)  
*elderi* Stirling and Zietz, 1893 (*mahoodi* Wells and  
 Wellington, 1985)  
*intermedius* Ogilby, 1892  
*jeanae* Storr, 1988  
*mcmillani* Storr, 1978  
*michaelseni* Werner, 1910  
*rankini* Storr, 1979  
*spinigerus* Gray, 1842 (*inornatus*\* Storr, 1988)

- strophurus* Duméril and Bibron, 1836 (*dumerilii* Fitzinger, 1843)  
*taeniatus* Lönnberg and Andersson, 1913  
*wellingtonae* Storr, 1988  
*williamsi* Kluge, 1963  
*wilsoni* Storr, 1983
- Tarentola* Gray, 1825 (*Makariogecko* Joger, 1984; *Sahelogecko* Joger, 1984; *Saharogecko* Joger, 1984; *Neotarentola* Joger 1984)  
*americana* Gray, 1831 (*milberti* Duméril and Bibron, 1836; *cubana* Gundlach and W. Peters, 1865; *warreni*\* Schwartz, 1968)  
*angustimentalis* Steindachner, 1891  
*annularis* I. Geoffroy, 1827 (*savignyi* Audouin, 1827; *aegyptiacus* Cuvier, 1829; *senegalensis* Boulenger, 1885; *quadraticauda* Toriner, 1905; *relicta*\* Joger, 1984)  
*boehmei* Joger, 1984  
*boettgeri* Steindachner, 1891 (*hierrensis*\* Joger and Bischoff, 1983; *bischoffi*\* Joger, 1984)  
*borneensis* Gray, 1845 (*gigas*\* Barboza du Bocage, 1875; *rudis*\* Boulenger, 1906; *protogigas*\* Joger, 1984; *maioensis*\* Schleich, 1984; *brancoensis* Schleich, 1984)  
*caboverdianus* Schleich, 1984 (*substituta*\* Schleich, 1984; *antoensis* Joger, 1984; *razianus*\* Schleich, 1984; *nicolauensis*\* Schleich, 1984)  
*clypeata* [?] Gray, 1842  
*darwini* Joger, 1984  
*delalandii* Duméril and Bibron, 1836  
*deserti* Boulenger, 1891  
*ephippiata* O'Shaughnessy, 1875 (*hoggarensis*\* Werner, 1937; *panousei* Pasteur, 1959; *senegambiae*\* Joger, 1984; *nikolausi*\* Joger, 1984)  
*gomerensis* Joger and Bischoff, 1983  
*mauritanica* Linnaeus, 1758 (*muricatus* Laurenti, 1768; *fascicularis*\* Daudin, 1802; *stellio* Merrem, 1820; *muralis* Duméril and Bibron, 1836; *facetanus* Strauch, 1862; *atlantica* Doumergue, 1899; *gracilis* Doumergue, 1899; *lissoide* Doumergue, 1899; *saharae* Doumergue, 1899; *tuberculata* Rosen, 1905; *juliae*\* Joger, 1984)  
*neglecta* Strauch, 1887 (*angusticeps* Strauch, 1887; *greyi*\* Joger, 1984)  
*parvicarinata* Joger, 1980
- Tenuidactylus* Shcherbak and Golubev, 1984  
*caspius* Eichwald, 1831 (*fasciatus* Menetries, 1832;

*insularis* Akhmedow and Shcherbak, 1978)  
*fedtschenkoi* Strauch, 1887  
*longipes* Nikolsky, 1896 (*microlepis*\* Lantz, 1918)  
*turcmenicus* Shcherbak, 1978  
*voraginosus* Leviton and S. Anderson, 1984

*Teratolepis* Günther, 1870  
*albofasciatus* Grandison and Soman, 1963  
*fasciata* Blyth, 1853

*Teratoscincus* Strauch, 1863  
*bedriagai* Nikolsky, 1899  
*microlepis* Nikolsky, 1899  
*przewalskii* Strauch, 1887  
*scincus* Schlegel, 1858 (*keyserlingii*\* Strauch,  
 1863; *zarudnyi* Nikolsky, 1896; *roborowskii*  
 Bedriaga, 1905; *rustamowi*\* Shcherbak, 1979)

*Thecadactylus* Oken, 1817 (*Thecodactylus* Wagler, 1830)  
*rapicauda* Houttuyn, 1782 (*perfoliatus* Schneider,  
 1793; *laevis* Daudin, 1802; *surinamensis*  
 Daudin, 1802; *theconyx* Duméril and Bibron,  
 1836; *tristis* Hallowell, 1854)

*Tropiocolotes* W. Peters, 1880  
*persicus* Nikolsky, 1903  
*scorteccii* Cherchi and Spano, 1963  
*steudneri* W. Peters, 1869 (*petersii* Boulenger,  
 1891; *natterei* Steindachner, 1900)  
*tripolitanus* W. Peters, 1880 (*occidentalis*\*  
 Parker, 1942; *somalicus*\* Parker, 1942;  
*algericus*\* Loveridge, 1947; *apoklomax*  
 Papenfuss, 1970)

*Urocotyledon* Kluge, 1983  
*inexpectata* Stejneger, 1893  
*palmata* Mocquard, 1902  
*weileri* L. Müller, 1909  
*wolterstorffi* Tornier, 1900

*Uroplatus* A. Duméril, 1805 [1806] (*Uroplates* Gray,  
 1825; *Rhacoessa* Wagler, 1830; *Oiacurus*  
 Fitzinger, 1843; *Lonchurus* Fitzinger, 1843)  
*alluaudi* Mocquard, 1894  
*ebenau* Boettger, 1879 (*boettgeri* Fischer, 1883  
 [1884]; *phantasticus* Boulenger, 1888;  
*schneideri* Lamberton, 1913)  
*fimbriatus* Schneider, 1797 (*hypoxantha* Wagler,  
 1833)  
*guentheri* Mocquard, 1908  
*lineatus* Duméril and Bibron, 1836  
*sikorae* Boettger, 1913